

Jeremy P. Watkins, M.D.

John B. McIntyre, M.D.

Sean M. Callahan, M.D.

5751 Edwards Ranch Rd. Fort Worth Tx 76109 Phone: 817-332-8848 Fax: 817-335-2670

www.fortworthent.net

TONSILLECTOMY/ADENOIDECTOMY

POST OPERATIVE INSTRUCTONS:

- Rest for 1-2 days. Resume normal activity as soon as possible. No heavy activity for 2 weeks
- Pain medication prescription will be given the day of surgery. Take pain medication every 4-6 hours with food. It is important to keep on a schedule with this medication to avoid gaps in coverage of pain relief.
- Swish, gargle and spit with prescribed mouthwash. Take every 4 hours, as needed.
- Drink plenty of fluid in small sips, as often as possible.
- Return to clinic1-2 weeks after surgery for follow up.

WHAT TO EXPECT:

Appearance of Tonsil Bed	Normal, white patches
Bleeding	If there is excessive bright red or brown drainage, gargle with ICE water (NO SALT WATER) for 10-20 minutes. If it doesn't stop, please call the office.
Pain	Can be severe with difficulty swallowing. Drinking plenty of liquids, taking the pain medication and resuming a normal diet will help. Days 6-10 is when the pain is at the highest level.
Voice	There may be a change in voice secondary to increased flow through the nose, this is usually transient.
Breath	Many have some bad breath for one week. Please use regular toothpaste only. DO NOT use regular mouth wash or oral rinses; only the prescribed mouthwash. Mints and gums may be helpful.
Nausea	Can last for 24 hours.
Temperature	Mild elevations for 48 hours.

EAR PAIN IS NORMAL.

Updated 4/25/2017